

Compound Formulation Active Pharmaceutical Ingredients and the Associated Adverse Effects and Potential Contraindications/Warnings and Precautions

## Moxi+Brom

## **Moxifloxacin HCl Possible Adverse Reaction Potential Contraindications / Warnings and Precautions** Conjunctivitis, Decreased visual acuity, Dry eye, Keratitis, Ocular Patients less than 1 year, Lactation, Pregnancy. Known discomfort, Ocular hyperemia, Ocular pain, Ocular pruritus, hypersensitivity to flouroquinolones or any of the components. Subconjunctival hemorrhage, and Tearing, Anaphylaxis, Growth Growth of resistant organisms with prolonged use. of resistant organisms, Fever, Increased cough, Infection, Otitis media, Pharyngitis, Rash and Rhinitis. **Bromfenac** Possible Adverse Reaction **Potential Contraindications / Warnings and Precautions** Anterior chamber eye inflammation, Headache, Vitrious Lactation, Pediatric patients, Pregnancy, Slow or Delayed Healing (NSAIDS), Potential for cross-sensitivity (NSAIDS) floaters, Iritis, Eye pain, Ocular hypertension, Burning sensation, Conjunctival hyperemia, Corneal erosion, Corneal or any of the components. Increased bleeding time of ocular perforation, Corneal thinning, Corneal ulcer, Epithelial tissue, Keratitis, Corneal reactions. Corneal adverse events keratopathy, Eye irritation, Eye pruritus, Eye redness, are increased in patients with dry eye syndrome, rheumatoid Hypersensitivity reaction, Keratitis, Prolonged bleeding, arthritis, repeat ocular surgeries in a short time, and diabetes Stinging sensation, Abnormal sensation in eyes, Anaphylaxis mellitus.

